

Doctor Discussion Guide



What's Your Sleep Story?

This doctor discussion guide can help you understand the many ways your difficulty with falling asleep or staying asleep (insomnia) is impacting your life.

Simply select the choice(s) that best fill in the blank. You can then share the results with your doctor to have a more empowered conversation about whether prescription QUVIVIQ is right for you.

I have trouble with select all that apply
FALLING ASLEEP STAYING ASLEEP WAKING UP TOO EARLY NOT FEELING WELL RESTED DURING THE DA
My insomnia occurs
NIGHTLY VERY OFTEN OCCASIONALLY ONCE A WEEK ONCE A MONTH RARELY
And I'm
SHRUGGING IT OFF RESIGNED TO IT FRUSTRATED AT MY LIMIT
I get about hours of sleep nightly.
0 TO 2 3 TO 4 5 TO 6 7 OR MORE
I have less daytime energy for select all that apply
RELATIONSHIPS SELF-CARE HOBBIES ERRANDS EXERCISING WORK/SCHOOL
I have tried, but I'm still looking for better results. select all that apply
PRACTICING HEALTHY SLEEP HABITS OTC SLEEP AID RX MEDICATION MELATONIN MEDITATION
Remember, bring this with you at your next appointment so your doctor can get a better sense of your sleep struggles and the two of you can discuss how QUVIVIQ may be able to help.

What is QUVIVIQ (daridorexant)?

QUVIVIQ is a prescription medicine for adults who have trouble falling asleep or staying asleep (insomnia).

Important Safety Information

Do not take QUVIVIQ if you fall asleep often at unexpected times (narcolepsy).

Please see additional Important Safety Information on pages 2-3. See full Prescribing Information and Medication Guide.



Doctor Discussion Guide



Insomnia impacts both your night and day.

QUVIVIQ is clinically proven to not only help you fall asleep faster, but stay asleep longer.

With more sleep, you may feel less tired during the day.

QUVIVIQ can help you:






Receive your first 30-day supply of QUVIVIQ for as little as \$0 and refills for as little as \$25* with the QSavings Card

*For eligible commercially insured patients. Terms and conditions apply.





For information on copay savings, support when taking QUVIVIQ, sleep tips, and more, register for Q&You today!

Important Safety Information

QUVIVIQ may cause serious side effects, including:

- Decreased awareness and alertness. The morning after you take QUVIVIQ, your ability to drive safely and think clearly may be decreased. You may also have sleepiness during the day.
- Do not take more QUVIVIQ than prescribed.
- Do not take QUVIVIQ unless you are able to stay in bed for at least 7 hours before you must be active again.
- Take QUVIVIQ at night within 30 minutes before going to bed.

Important Safety Information

QUVIVIQ is a federally controlled substance because it can be abused or lead to dependence.

Before taking QUVIVIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of depression, mental illness, or suicidal thoughts or actions; drug or alcohol abuse or addiction; a sudden onset of muscle weakness (cataplexy); daytime sleepiness
- have lung or breathing problems, including sleep apnea
- have liver problems
- are pregnant or plan to become pregnant
- are breastfeeding or plan to breastfeed

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins,

and herbal supplements.

- Taking QUVIVIQ with certain medicines can cause serious side effects. QUVIVIQ may affect the way other medicines
 work and other medicines may affect the way QUVIVIQ works.
- · Do not take QUVIVIQ with other medicines that can make you sleepy unless instructed by your healthcare provider.

What should I avoid while taking QUVIVIQ?

- · Do not drink alcohol while taking QUVIVIQ. It can increase the effects of alcohol, which can be dangerous.
- Do not drive, operate heavy machinery, do anything dangerous, or do other activities that require clear thinking if you do not feel fully awake, or you have taken QUVIVIQ and have less than a full night of sleep (at least 7 hours), or if you have taken more QUVIVIQ than prescribed.

QUVIVIQ may cause other serious side effects, including:

- Worsening depression and suicidal thoughts. Call your healthcare provider right away if you have any worsening depression or thoughts of suicide or dying.
- Temporary inability to move or talk (sleep paralysis) for up to several minutes, or hallucinations while you are going to sleep or waking up.
- **Complex sleep behaviors** such as sleep-walking, sleep-driving, preparing and eating food, making phone calls, having sex or doing other activities while not fully awake that you may not remember the next morning. Stop taking QUVIVIQ and call your healthcare provider right away if you experience a complex sleep behavior.

The most common side effects of QUVIVIQ are headache and sleepiness.

These are not the only side effects of QUVIVIQ. Call your doctor for advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information and Medication Guide.



Visit QUVIVIQ.com for more information

